

BAD BUGS:

SOME BUGS USE DIFFERENT METHODS TO DAMAGE VEGETABLES, FRUIT, SHRUBS OR TREES:

SAP SUCKERS: aphids, mites, thrips, mealy bugs, leafhoppers

LEAF LOVERS: hornworms, beetles, squash bugs, cabbage loopers

FRUIT FEEDERS: fruit flies, hornworms, plant bugs, yellow jackets,
hornets

ROOT WRECKERS: slugs, cutworms, nematodes, cabbage and onion
maggots, weevil larvae, white grubs



Stink Bug



Agave Weevil



Aphid



Armyworm



Asparagus Beetle



Bagworm



Bark Beetle



Boxelder Bug



Bark Louse



Blister Beetle



Cabbage Looper



Cankerworm



Chinch Bug



Cicada



Click Beetle



Codling Moth



Colorado Potato Beetle



Corn Earworm



Cottony-Cushion Scale



Cricket



Cucumber Beetle



Cutworm



Flea Beetle



Fleahopper



Fungus Gnat



Grape Leaf Skeletonizer Western



Grasshopper



Gypsy Moth



Harlequin Bug



Hickory Shuckworm



Horse Fly



Imported Cabbageworm



June Beetle



Kissing Bug



Leaffooted Bug



Leafhopper



Leafminer



Leafroller



Locust Borer



Mealybug



Mexican Bean Beetle



Mosquito



Orange Tortrix



Peach Tree Borer



Phylloxera



Pine Tip Moth



Plum Curculio



Psyllid



Puss Caterpillar



Sawfly



Scale



Sharpshooter



Shot Hole Borer



Slug



Snail



Sod Webworm



Spider Mite



Spittlebug



Springtail



Squash Bug



Tarnished Plant Bug



Tent Caterpillar



Termite



Thrips



Tobacco Hornworm



Tomato Hornworm



Tomato Hornworm Moth



Tomato Pinworm



Tomato Pinworm Moth



Tortoise Beetle



Treehopper



Twig Girdler



Webworm



Weevil



Whitefly